

New Perspective of Multi-dimensional Approach for the Management of Attention-deficit Hyperactivity Disorder: A Review



Sachitanand Singh^{1,*} , Renu Thakur² , Krishan Kumar³ and Ayush Dogra⁴

¹Department of Optometry, UIAHS, Chandigarh University, Punjab, India

²Centre for Research Impact and Outcome, Chitkara University, Punjab, India

³Department of Psychiatry, Postgraduate Institute of Medical Education & Research (PGIMER), Chandigarh, India

⁴Chitkara University Institute of Engineering and Technology, Chitkara University, Punjab, India

Abstract:

One of the most common mental diseases in childhood, attention-deficit/hyperactivity disorder (ADHD) often lasts into adulthood for many individuals. The neurodevelopmental condition known as ADHD impacts three areas of the brain: hyperactivity, impulsivity, and attention. The visual field is where attention is most affected by ADHD. Non-strabismic binocular vision disorder (NSBVD), which impairs eye coordination and makes it challenging to focus, has been linked to ADHD. Through a critical cognitive process called visual attention, humans are able to take in and organize information from their visual environment. This greatly affects how one observes, processes, and understands visual information in day-to-day living. Vision therapy is a non-invasive therapeutic approach that aims to improve visual talents and address visual attention deficits. This study aims to provide an overview of the research on the many approaches to treating ADHD, the relationship between NSBVD and ADHD, and whether vision therapy is a viable treatment option for ADHD. After a comprehensive search of many online resources, relevant studies were found. The review's findings provide insight into the range of ADHD patients' treatment choices. In order to improve treatment outcomes, non-pharmacological treatments can be employed either alone or in conjunction with medicine. Medicine by itself is insufficient and has several severe side effects when used continuously. The efficacy of vision therapy in improving visual attention and making recommendations for potential directions for further research in this field. Multiple studies are needed to identify the most effective treatment modalities for achieving positive outcomes for ADHD patients.

Keywords: Keywords, Attention-deficit/hyperactivity disorder (ADHD), Non-strabismic binocular vision disorder (NSBVD), Attention deficit disorder (ADD), Vision Therapy, Visualattention, Attention, Multidisciplinary ADHD management.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to this author at the Department of Optometry, UIAHS, Chandigarh University, Punjab, India; Tel: 7411949632; E-mails: Sachi.publication@gmail.com, Snsingh.social@gmail.com and Sachitanand.singh@chitkara.edu.in

Cite as: Singh S, Thakur R, Kumar K, Dogra A. New Perspective of Multi-dimensional Approach for the Management of Attention-deficit Hyperactivity Disorder: A Review. Open Neuroimaging J, 2024; 17: e18744400326671. <http://dx.doi.org/10.2174/0118744400326671241002074115>



Received: July 24, 2024
Revised: September 12, 2024
Accepted: September 18, 2024
Published: October 10, 2024



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Attention involves the cognitive mechanism of deliberately focusing on particular elements within the environment while disregarding others. It allows individuals to allocate their limited cognitive resources to relevant information, facilitating perception, memory,

decision-making, and action [1]. Visual attention is a cognitive process through which humans selectively focus on specific visual stimuli while filtering out irrelevant information. It plays a crucial role in perception, guiding gaze, and allocating cognitive resources to relevant visual elements in our environment [2, 3]. This phenomenon is closely tied to the mechanisms of visual perception, visual

search, and object recognition [4, 5]. Understanding visual attention has broad implications for various fields, including psychology, neuroscience, computer vision, and artificial intelligence. Selective attention, divided attention, and sustained attention are some of the different components and processes that make up visual attention. The ability to focus on specific stimuli while filtering out competing distractions was known as selective attention. On the other hand, divided attention entails efficiently allocating attention among numerous inputs or tasks at once [6, 7]. This filtration process was a part of visual information processing where all information reaches a different cognitive level, and whichever place our mind wants to see, that information reaches the highest level of thinking, and people would give attention to that object [8].

1.1. Visual Information Processing

The human visual system is bombarded with an overwhelming amount of visual information processing at any given moment. However, our cognitive resources are limited, and it is not feasible to process all incoming visual stimuli simultaneously. Therefore, visual attention acts as a mechanism that prioritizes relevant information while suppressing or ignoring irrelevant or distracting elements, as shown in Fig. (1). The visual information process was broadly divided into three subsections: visual-spatial,

visual analysis, and visual-motor. Visual analysis was further divided into four subsections, as mentioned in Fig. (2) [10-13].

1.2. Attention-deficit/hyperactivity Disorder (ADHD)

Attention, impulsivity, and hyperactivity are all impacted by the neurodevelopmental conditions called ADHD and ADD. Historically, ADD was used to describe the subtype of ADHD. However, in the current diagnostic criteria, According to the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), which was shown in Tables 1-2 [14, 15], ADHD is now the umbrella term used to describe both the hyperactive/impulsive and inattentive subtypes and ADD is no longer used as a separate diagnosis. Both children and adults can be affected by ADHD, which has a substantial negative influence on a number of facets of a person's life, including social relationships, work productivity, and academic achievement [16-18]. ADHD is thought to result from a confluence of genetic, environmental, and neurological factors, however, its specific causes were yet unknown, but a few conditions were directly related to ADHD, like neurotransmitter imbalance, maternal smoking and substance abuse, premature birth, low birth weight, and socioeconomic conditions [19]. Understanding ADHD and its complexities was crucial for effective diagnosis, treatment, and support for those affected by this disorder.




Fig. (1). This figure was adapted from Broadbent's Filter Model for Attention (1958) [9] and shows how the multiple unwanted impulses get filtered out and allow only the required information to the brain. In the Figure 'A' and 'B' are input, and with the help of the sensory pathway, reach selective filtration level where brain understands the information related to objects shape, size, colour and many more and then select only one information that was high-level processing and allow only 'A' information, and the memory also recognized the information 'A' [9-11].




Fig. (2). This figure was adapted from the visual perception book. This Figure shows how visual information processing was divided into visual-spatial, visual analysis, and visual motor for recognition of any objects, and it also shows that visual analyzing was done based on perceptual attention memory visualization.

The prevalence of ADHD varies across different populations and is influenced by various factors, including diagnostic criteria, assessment methods, cultural differences, and sample characteristics [15, 20, 21]. The reported prevalence rates also differ between children and adults. According to the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5), which was widely used for diagnosing ADHD, the prevalence of ADHD in children was approximately 5-10% and 2-5% [15] in adults worldwide. However, prevalence rates can differ among countries. In India, the prevalence rate was 1.30%

to 28.9% [20] in children and adolescents, and in north Indian schoolchildren, ADHD prevalence is 6.34% [21]. The attention span decreased after COVID-19 by 54.4% in students, which caused difficulty in academic performance [22]. With appropriate support and interventions, individuals with ADHD can develop strategies to cope with their symptoms, improve their focus and attention, and lead fulfilling lives. During comprehensive eye testing of ADHD symptomatic patients, it was observed that they had some form of binocular vision abnormality like convergence insufficiency 30%, accommodative dys-

function 24%, accommodative with vergence dysfunction 13%, and fusional vergence dysfunction 3%. Seventy-one % of the students reported positive ADHD symptoms, and the majority of the students exhibited inattention issues rather than hyperactivity [23]. Fifty-two% of the kids had non-strabismic binocular vision disorder (NSBVD) and ADHD symptoms. Home-based visual treatment alleviated the attention deficit disorder [24, 25]. NSBVD abnormality refers to an eye condition characterized by an impairment in the coordination and alignment of the two eyes. People with attention deficit hyperactivity disorder (ADHD) have been reported to have NSBVD [23]. In individuals with non-strabismic binocular vision abnormality, the eyes struggle to work together efficiently, leading to difficulties in focusing, tracking moving objects, eye strain, double vision, reduced attention and reading comprehension, and perceiving depth and spatial relationships. Research has shown a higher prevalence of binocular vision abnormalities, such as convergence insufficiency, divergence excess, and accommodative dysfunction, among individuals with ADHD [23, 24, 26]. The exact relationship between ADHD and non-strabismic binocular vision abnormalities is still being explored. It is believed that certain neurological and developmental factors like depression, anxiety, and autism spectrum disorder (ASD) associated with ADHD may contribute to the development or exacerbation of these visual abnormalities [27, 28]. Early detection and diagnosis of non-strabismic binocular vision abnormalities in individuals with ADHD are crucial for appropriate intervention [29]. Visual attention can be improved with the help of vision therapy. Vision therapy is a type of rehabilitation program that involves exercises and activities aimed at improving visual skills and abilities.

These skills include eye movement control, focusing abilities, binocular vision, visual processing skills, and attention. The goal of vision therapy is to improve overall visual function, reduce symptoms related to visual dysfunction, and enhance the individual's quality of life [26]. A new technology, which was based on electroencephalography (EEG), was a non-invasive technique to diagnose, measure, and help in vision therapy. The electrical activity of the brain is electroencephalography (EEG), which is aided by a brain-computer interface (BCI). It assists in the diagnosis and management of ADHD patients [30-32].

2. METHODOLOGY

All the articles selected were related to these keywords: "attention deficit disorder" OR "ADD" OR "attention deficit hyperactive symptoms" OR "ADHS" OR "attention deficit hyperactive disorder" OR "ADHD" AND "binocular vision abnormality" OR "non-strabismic binocular vision dysfunction" OR "NSBVD" OR "accommodation" OR "convergence" AND "ADHD Treatment" OR "ADHD Diagnosis." Seven hundred eighty articles were found in PubMed, Scopus, Web of Science, and Google Scholar search engines between 2014 and 2023. A total of 57 research articles were considered for this review after filtering out the review articles, conference proceedings, and book chapters. The articles about ADHD and its diagnosis and management were considered for this review, as shown in Fig. (3). The main objectives of this review were to find out the different treatment modalities available for ADHD, the relationship between ADHD and non-strabismic binocular vision dysfunction, and the impact of vision therapy as a treatment option for ADHD.

Table 1. Key aspects of the assessment and diagnosis process for ADHD.

<ul style="list-style-type: none"> • Clinical Interviews • Symptom Questionnaires [38-40] like Adult ADHD self-report scale V1.1(ASRS-V1.1) and Adult self-report screening scale for DMS-5(ASRS-5) • Medical and Psychological Evaluation • Behavioral observations • Psychological Testing • Diagnostic Criteria Diagnostic and Statistical Manual of Mental Disorders (DSM-5) • Comprehensive eye examinations
--

Table 2. The diagnostic and statistical manual of mental disorders, fifth edition (DSM-5), published by the american psychiatric association, provides the following diagnostic criteria for ADHD [14, 15, 19].

<p>Inattention: Six or more symptoms of inattention have persisted for at least six months to a degree that is inconsistent with the developmental level:</p> <ul style="list-style-type: none"> ■ Pays insufficient attention to details or commits thoughtless errors at work, school, or in other activities. ■ Has trouble maintaining focus during tasks or play activities. ■ Does not appear to pay attention when directly addressed. ■ Ignores directions and does not finish their homework, housework, or work-related responsibilities. ■ Has trouble planning activities and tasks. ■ Is unwilling to perform tasks that demand prolonged mental effort because they are unpleasant or uncomfortable. ■ Misplaces items required for jobs or activities. ■ Easily sidetracked by unrelated stimuli. ■ Forgetful throughout routine tasks. 	<p>Hyperactivity and Impulsivity: Six or more symptoms of hyperactivity-impulsivity have persisted for at least six months to a degree that is inconsistent with the developmental level:</p> <ul style="list-style-type: none"> ■ Moves their hands, feet, or seat while fidgeting. ■ Stands up while it is expected that they stay seated. ■ Moves around or climbs in inappropriate places (in adults or teenagers, may be restricted to irrational emotions of restlessness). ■ Unable to play or partake in quiet activities. ■ "On the go," acting as if "driven by a motor." ■ Talks way too much. ■ Answers are blurted out before all the questions have been answered. ■ Has trouble waiting for their turn. ■ Intrudes on or interrupts others.
--	--




Fig. (3). Paper selection flow chart. The flowchart provides information about the studies that were ultimately included in this review, as well as the search approach.

4. RESULTS

4.1. COMORBIDITIES AND ASSOCIATED CONDITIONS

Comorbidities and associated conditions can significantly impact the lives of individuals with ADHD, exacerbating the challenges they face and requiring additional support and treatment. Some of the commonly observed comorbidities and associated conditions of ADHD, such as, oppositional defiant disorder (ODD) and conduct disorder (CD), learning disabilities, mood disorders, autism spectrum disorder (ASD), substance abuse disorders, sleep disorders, tic disorders, and obesity have been discussed in this study. It is important to note that not all individuals with ADHD experience comorbidities or associated conditions. The presence of these conditions varies among individuals, and the severity can also differ. It is important to note that with appropriate interventions and support, children with ADHD can thrive and lead fulfilling lives. Adults with ADHD tend to experience more internal restlessness rather than external hyperactivity. Living with ADHD as an adult can present numerous challenges. They may find it challenging to meet deadlines, prioritize tasks, and sustain attention during lengthy or monotonous activities. This can lead to increased stress, underachievement, feelings of frustration, and low self-esteem. Adults with ADHD may also face challenges in their personal

relationships [29, 33]. Assessment and diagnosis of Attention-Deficit/Hyperactivity Disorder (ADHD) was a big challenge to depreciate from its comorbidity. ADHD diagnosis involves a comprehensive evaluation of an individual's symptoms, behavior, and history to determine the presence and severity of the disorder [34]. The assessment process typically includes multiple sources of information, such as interviews, questionnaires, observations, and psychological testing, along with comprehensive eye examinations to assess binocular vision function. A common onset for ADHD occurs in childhood, and it can last throughout adolescence and maturity. Before the age of 12, several symptoms were evident and can be seen in two or more contexts (such as the home, school, workplace, or social setting). It is important to note that only qualified healthcare professionals, such as psychiatrists, psychologists, optometrists, or pediatricians, should perform the assessment and diagnosis of ADHD [35-37].

It is important to note that diagnosing ADHD is a complex process, and misdiagnosis or overdiagnosis can occur. A comprehensive assessment ensures that other possible explanations for the symptoms are considered and that an accurate diagnosis is made. Overall, the assessment and diagnosis of ADHD involve a multifaceted approach [6, 35] that integrates information from various sources, as shown in Table 1.

4.2. Multimodal Management

Managing ADHD typically involves a multimodal treatment approach that combines different strategies and interventions tailored to the individual's specific needs. This approach recognizes the heterogeneity of ADHD symptoms and acknowledges that a "one-size-fits-all" approach may not be effective for everyone. What works for one individual may not work for another, therefore, a personalized, complementary, and alternative medicine approach is essential [41]. It typically involves a combination of medication, behavioral interventions, psychoeducation, parent-child interaction therapy, and support [42]. Medication is often a cornerstone of ADHD treatment, especially for individuals with moderate to severe symptoms. Stimulant medications, such as methylphenidate or amphetamines, are commonly prescribed [43] and have been shown to reduce hyperactivity and impulsivity and improve attention and executive functioning in many individuals, and all the

treatment options are available in Fig. (4). However, medication alone may not be sufficient for comprehensive ADHD management [44], and long uses of medication create multiple adverse effects on children, adolescents, and adults, such as decreased appetite, weight loss, dyspepsia, abdominal pain, stomach ache, irritability, mood disorders, dizziness, anorexia, nausea, somnolence, and vomiting [45-47]. Non-pharmacological interventions play a crucial role in the treatment of attention-deficit/hyperactivity disorder (ADHD). Non-pharmacological approaches can be used as standalone interventions or in combination with medication to enhance the overall treatment outcome. These interventions focus on addressing various aspects of ADHD, such as improving attention, impulsivity control, organizational skills, and behavioral management [48]. Some commonly employed non-pharmacological treatment approaches for ADHD are given in the study [43, 49, 50], including electroencephalographic (EEG) based, which are used for diagnostics and management purposes both [51-61].




Fig. (4). Multidisciplinary ADHD treatment options available, including pharmacological and non-pharmacological.

5. DISCUSSION

Vision therapy for NSBVD involves a series of specialized eye exercises and visual activities tailored to the individual's specific needs, and it is also dependent on individual refractive status [62], vergence [63], stereo acuity, fusional vergence amplitudes [64], convergence insufficiency [25, 65, 66], near point of convergence (NPC) [24], accommodation problem [67], saccade [68] This condition could be improved by the help of vision therapy or orthoptic exercise. The treatment is typically conducted under the guidance of a trained optometrist or, vision therapist, or orthoptist and follows a structured program. The therapy aims to enhance the coordination and teamwork between the eyes, improve eye movement control, enhance focusing abilities, and strengthen visual processing skills, which is shown in Fig. (2) [69-73].

Table 3. The specific techniques used in vision therapy for NSBVD may include [69-73, 73].

- Eye teaming exercises
- Visual tracking exercises
- Visual perception activities
- Accommodative exercises
- Vergence exercises
- Visual-motor integration tasks

Furthermore, by addressing NSBVD through vision therapy techniques, which were mentioned in Table 3, ADHD patients may experience several benefits. Improved eye coordination and focusing abilities can positively impact reading skills, attention, and concentration. Enhanced visual processing and perception skills can contribute to better comprehension, visual memory, and overall academic performance [74]. Reducing visual discomfort and fatigue can alleviate the strain on the visual system, potentially leading to a reduction in ADHD-related symptoms. It is significant to stress that vision therapy should be incorporated as part of a thorough treatment strategy for those with ADHD [74]. Consulting with healthcare professionals, such as psychologists, occupational therapists, optometrists, and educators, can help determine the most suitable treatment plan and ensure a holistic approach to address the various aspects of the condition for each individual. There were a few studies that showed that signal processing and convolution neural networks can help to diagnose and management of brain-related problems [75-78]]. Furthermore, the evolving understanding of ADHD through comprehensive research, clinicians and partitioners can improve diagnostic accuracy, develop targeted treatments, and enhance the overall well-being and quality of life for individuals with ADHD. While medication and behavioral therapies are commonly used in the treatment of ADHD, researchers and clinicians continue to explore alternative approaches to improve outcomes for individuals with this condition. The possible involvement of vision therapy treatment is one area of investigation, particularly in resolving non-strabismic binocular vision impairments in ADHD patients [79].

CONCLUSION

The current treatment strategy for ADHD includes a combination of medication, behavioral interventions, psychotherapy, occupational therapy, psychoeducation, parent-child interaction therapy, and support, which has shown improvement, but this treatment does not include vision-related treatment. Several studies show that non-strabismic binocular vision dysfunction (NSBVD) has been linked to ADHD symptoms, and vision therapy for NSBVD improved ADHD symptoms, although there was still more research needed. The potential advantages of vision therapy designed especially for people with ADHD and non-strabismic binocular vision impairments have only been briefly studied. There is currently a research gap in determining the efficacy of vision therapy as a treatment option for ADHD symptoms despite the intriguing association between non-strabismic binocular vision impairments and ADHD. Future studies should concentrate on carrying out sizable, randomized controlled trials that assess the effects of vision therapy on ADHD symptoms in people with non-strabismic binocular vision impairments in order to fill this knowledge gap. Moreover, by closing the research gap in the field of vision therapy for non-strabismic binocular vision abnormalities in ADHD, clinicians and practitioners can gain a deeper understanding of the potential benefits and limitations of this treatment approach. Such knowledge can guide clinicians in developing comprehensive and personalized interventions that consider the unique visual needs of individuals with ADHD, leading to improved outcomes and quality of life for those affected by this condition.

AUTHOR'S CONTRIBUTION

Study conception and design: Renu Thakur, Krishan Kumar, Ayush Dogra

Data collection: Sachitanand Singh

Analysis and interpretation of results: Sachitanand Singh, Renu Thakur, Krishan Kumar

Draft manuscript: Sachitanand Singh

Review of Final Draft of Manuscript: Sachitanand Singh, Renu Thakur, Krishan Kumar, Ayush Dogra

CONSENT FOR PUBLICATION

Not applicable.

FUNDING

None.

CONFLICT OF INTEREST

There were no conflicts of interest financial or otherwise.

ACKNOWLEDGEMENTS

Thanks to Dr. Sonika Bakshi, Dean of the Chitkara School of Health Sciences at Chitkara University, Punjab, and Dr. Archana Mantri, Vice Chancellor of Chitkara University, Punjab.

REFERENCES

- [1] Broman AT, West SK, Muñoz B, Bandeen-Roche K, Rubin GS, Turano KA. Divided visual attention as a predictor of bumping while walking: The Salisbury eye evaluation. *Invest Ophthalmol Vis Sci* 2004; 45(9): 2955-60.
<http://dx.doi.org/10.1167/iovs.04-0219> PMID: 15326107
- [2] Binder MD, Hirokawa N, Windhorst U. *Encyclopedia of neuroscience*. Berlin, Germany: Springer 2009; p. 3166.
<http://dx.doi.org/10.1007/978-3-540-29678-2>
- [3] Kastner S, McMains SA, Beck DM. Mechanisms of selective attention in the human visual system: Evidence from neuroimaging. 2019; (4): 205-17.
- [4] Ball K, Owsley C, Sloane ME, Roenker DL, Bruni JR. Visual attention problems as a predictor of vehicle crashes in older drivers. *Invest Ophthalmol Vis Sci* 1993; 34(11): 3110-23. PMID: 8407219
- [5] Thakur R, Jayakumar J, Pant S. A comparative study of visual attention in hearing impaired and normal schoolgoing children. *Indian Journal of Otolaryngology* 2019; 25(4): 3-5.
http://dx.doi.org/10.4103/indianjotol.INDIANJOTOL_19_19
- [6] Zagaykan Y, Spryn O, Zagaykan N. Research Of sensorimotor reaction, memory and attention indices under sensory deprivation. *EUREKA: Life Sciences* 2019; (5): 3-12.
- [7] Biederman J. Attention-deficit/hyperactivity disorder: A selective overview. *Biol Psychiatry* 2005; 57(11): 1215-20.
<http://dx.doi.org/10.1016/j.biopsych.2004.10.020> PMID: 15949990
- [8] Müller HJ, Krummenacher J. Visual search and selective attention. *Vis Cogn* 2006; 14(4-8): 389-410.
<http://dx.doi.org/10.1080/13506280500527676>
- [9] Sullivan L. Selective attention and secondary message analysis: A reconsideration of Broadbent's filter model of selective attention. *Q J Exp Psychol* 1976; 28(2): 167-78.
<http://dx.doi.org/10.1080/14640747608400549>
- [10] Buchholz J, Aimola Davies A. Adults with dyslexia demonstrate attentional orienting deficits. *Dyslexia* 2008; 14(4): 247-70.
<http://dx.doi.org/10.1002/dys.356> PMID: 18023001
- [11] Desimone R, Duncan J. Neural mechanisms of selective visual attention. *Annu Rev Neurosci* 1995; 18(1): 193-222.
<http://dx.doi.org/10.1146/annurev.ne.18.030195.001205> PMID: 7605061
- [12] Deutsch JA, Deutsch D. Attention: Some theoretical considerations. *Psychol Rev* 1963; 70(1): 80-90.
<http://dx.doi.org/10.1037/h0039515> PMID: 14027390
- [13] Lund N. *Attention and pattern recognition*. London: Routledge 2020.
<http://dx.doi.org/10.4324/9781003060185>
- [14] Epstein JN, Loren REA. Changes in the definition of ADHD in DSM-5: Subtle but important. *Neuropsychiatry (London)* 2013; 3(5): 455-8.
<http://dx.doi.org/10.2217/npj.13.59> PMID: 24644516
- [15] American Psychiatric Association. *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, DC: American Psychiatric Publishing 2013.
- [16] Kirk HE, Gray K, Riby DM, Taffe J, Cornish KM. Visual attention and academic performance in children with developmental disabilities and behavioural attention deficits. *Dev Sci* 2017; 20(6): e12468.
<http://dx.doi.org/10.1111/desc.12468> PMID: 27649816
- [17] Lai SA, George Benjamin R, Schwanenflugel PJ, Kuhn MR. The longitudinal relationship between reading fluency and reading comprehension skills in second-grade children. *Read Writ Q* 2014; 30(2): 116-38.
<http://dx.doi.org/10.1080/10573569.2013.789785>
- [18] Wei CC, Ma MY. Influences of visual attention and reading time on children and adults. *Read Writ Q* 2017; 33(2): 97-108.
<http://dx.doi.org/10.1080/10573569.2015.1092100>
- [19] Martel MM, Schimmack U, Nikolas M, Nigg JT. Integration of symptom ratings from multiple informants in ADHD diagnosis: A psychometric model with clinical utility. *Psychol Assess* 2015; 27(3): 1060-71.
<http://dx.doi.org/10.1037/pas0000088> PMID: 25730162
- [20] Joseph J, Devu B. Prevalence of attention-deficit hyperactivity disorder in India: A systematic review and meta-analysis. *Indian J Psychiatr Nurs* 2019; 16(2): 118-25.
http://dx.doi.org/10.4103/IOPN.IOPN_31_19
- [21] Gupta RK, Sharma P, Banal R, et al. Prevalence and correlates of Attention Deficit Hyperactive Disorder (ADHD) risk factors among school children in a rural area of North India. *J Family Med Prim Care* 2020; 9(1): 115-8.
http://dx.doi.org/10.4103/jfmpc.jfmpc_587_19 PMID: 32110575
- [22] Quintiliani L, Sisto A, Vicinanza F, Curcio G, Tambone V. Resilience and psychological impact on Italian university students during COVID-19 pandemic. Distance learning and health. *Psychol Health Med* 2022; 27(1): 69-80.
<http://dx.doi.org/10.1080/13548506.2021.1891266> PMID: 33602027
- [23] Sharma S, Sarkar S. Attention deficit hyperactivity disorder (ADHD) symptoms among university students associated with non-strabismic binocular vision dysfunctions (NSBVDs). *Optom Vis Perform* 2021; 9(1): 30-8.
- [24] Dawidowsky B, Cerovski B, Klobučar A, Dawidowsky K. Do orthoptic exercises have any influence on children and adolescents diagnosed with convergence insufficiency and attention deficit/hyperactivity disorder? *Acta Clin Croat* 2019; 58(4): 662-71.
<http://dx.doi.org/10.20471/acc.2019.58.04.14>
- [25] Granet DB, Gomi CF, Ventura R, Miller-Scholte A. The relationship between convergence insufficiency and ADHD. *Strabismus* 2005; 13(4): 163-8.
<http://dx.doi.org/10.1080/09273970500455436> PMID: 16361187
- [26] Scheiman M, Wick B. *Clinical management of binocular vision: heterophoric, accommodative, and eye movement disorders*. Lippincott Williams & Wilkins 2008.
- [27] Howard IP, Rogers BJ. *Binocular vision and stereopsis*. Oxford University Press 1995.
- [28] Rommelse NNJ, Van der Stigchel S, Sergeant JA. A review on eye movement studies in childhood and adolescent psychiatry. *Brain Cogn* 2008; 68(3): 391-414.
<http://dx.doi.org/10.1016/j.bandc.2008.08.025> PMID: 18835079
- [29] Spencer TJ. ADHD and comorbidity in childhood. *J Clin Psychiatry* 2006; 67: 27-31. PMID: 16961427
- [30] Khullar V, Salgotra K, Singh HP, Sharma DP. Deep learning-based binary classification of ADHD using resting state MR images. *Augment Hum Res* 2021; 6(1): 5.
<http://dx.doi.org/10.1007/s41133-020-00042-y>
- [31] Henry JC. *Electroencephalography: Basic principles, clinical applications, and related fields*. Neurology 2006; 67(11): 2092.
<http://dx.doi.org/10.1212/01.wnl.0000243257.85592.9a>
- [32] Carr N. *The shallows: What the internet is doing to our brains*. WW Norton & Company 2020.
- [33] Katzman MA, Bilkey TS, Chokka PR, Fallu A, Klassen LJ. Adult ADHD and comorbid disorders: Clinical implications of a dimensional approach. *BMC Psychiatry* 2017; 17(1): 302.
<http://dx.doi.org/10.1186/s12888-017-1463-3> PMID: 28830387
- [34] Steinhausen H-C. The heterogeneity of causes and courses of attention-deficit/hyperactivity disorder. *Acta Psychiatr Scand* 2009; 120(5): 392-9.
<http://dx.doi.org/10.1111/j.1600-0447.2009.01446.x> PMID: 19807721
- [35] Zametkin AJ, Ernst M. Problems in the management of attention-deficit-hyperactivity disorder. *N Engl J Med* 1999; 340(1): 40-6.
<http://dx.doi.org/10.1056/NEJM199901073400107> PMID: 9878644
- [36] Biaggio MK, Bittner E. Psychology and optometry: Interaction and collaboration. *Am Psychol* 1990; 45(12): 1313-5.
<http://dx.doi.org/10.1037/0003-066X.45.12.1313.a> PMID: 2285180
- [37] Felt BT, Biermann B, Christner JG, Kochhar P, Harrison RV. Diagnosis and management of ADHD in children. *Am Fam Physician* 2014; 90(7): 456-64.

- PMID: 25369623
- [38] Silverstein MJ, Faraone SV, Alperin S, Biederman J, Spencer TJ, Adler LA. How informative are self-reported adult attention-deficit/hyperactivity disorder symptoms? An examination of the agreement between the adult attention-deficit/hyperactivity disorder self-report scale V1. 1 and adult attention-deficit/hyperactivity disorder investigator symptom rating scale. *J Child Adolesc Psychopharmacol* 2018; 28(5): 339-49. <http://dx.doi.org/10.1089/cap.2017.0082> PMID: 29172673
- [39] Kiatrungrit K, Putthirisi S, Hongsanguansri S, Wisajan P, Jullagate S. Validity and reliability of adult ADHD self-report scale Thai version (ASRS-V1. 1 TH). *Shanghai Jingshen Yixue* 2017; 29(4): 218-27. PMID: 28955141
- [40] Loskutova NY, Lutgen CB, Callen EF, Filippi MK, Robertson EA. Evaluating a web-based adult ADHD toolkit for primary care clinicians. *J Am Board Fam Med* 2021; 34(4): 741-52. <http://dx.doi.org/10.3122/jabfm.2021.04.200606> PMID: 34312267
- [41] Pellow J, Solomon EM, Barnard CN. Complementary and alternative medical therapies for children with attention-deficit/hyperactivity disorder (ADHD). *Altern Med Rev* 2011; 16(4): 323-37. PMID: 22214252
- [42] Baydala L, Wikman E. The efficacy of neurofeedback in the management of children with attention deficit/hyperactivity disorder. *Paediatr Child Health* 2001; 6(7): 451-5. <http://dx.doi.org/10.1093/pch/6.7.451> PMID: 20107553
- [43] Hesslinger B, Tebartz van Elst L, Nyberg E, *et al.* Psychotherapy of attention deficit hyperactivity disorder in adults. *Eur Arch Psychiatry Clin Neurosci* 2002; 252(4): 177-84. <http://dx.doi.org/10.1007/s00406-002-0379-0> PMID: 12242579
- [44] Behiyat S. Attention deficit hyperactivity disorder treatment review. *Hum Behav Dev Soc* 2011; 6(1): 12-7.
- [45] Wang Y, Zheng Y, Du Y, *et al.* Atomoxetine versus methylphenidate in paediatric outpatients with attention deficit hyperactivity disorder: A randomized, double-blind comparison trial. *Aust N Z J Psychiatry* 2007; 41(3): 222-30. <http://dx.doi.org/10.1080/00048670601057767> PMID: 17464703
- [46] Cortese S, Panei P, Arcieri R, *et al.* Safety of methylphenidate and atomoxetine in children with attention-deficit/hyperactivity disorder (ADHD): Data from the Italian National ADHD Registry. *CNS Drugs* 2015; 29(10): 865-77. <http://dx.doi.org/10.1007/s40263-015-0266-7> PMID: 26293742
- [47] Holmskov M, Storebø OJ, Moreira-Maia CR, *et al.* Gastrointestinal adverse events during methylphenidate treatment of children and adolescents with attention deficit hyperactivity disorder: A systematic review with meta-analysis and Trial Sequential Analysis of randomised clinical trials. *PLoS One* 2017; 12(6): e0178187. <http://dx.doi.org/10.1371/journal.pone.0178187> PMID: 28617801
- [48] Ramsay JR, Rostain AL. Cognitive behavioral therapy for adult ADHD: An integrative psychosocial and medical approach. Routledge 2014.
- [49] Tarver J, Daley D, Sayal K. Attention-deficit hyperactivity disorder (ADHD): An updated review of the essential facts. *Child Care Health Dev* 2014; 40(6): 762-74. <http://dx.doi.org/10.1111/cch.12139> PMID: 24725022
- [50] Maxwell V. Diagnosis and treatment of the gifted student with attention deficit disorder: A structure of intellect (SOI) approach. *J Read Writ Learn Disabil Int* 1989; 5(3): 247-52. <http://dx.doi.org/10.1080/0748763890050304>
- [51] Esqueda-Elizondo JJ, Juárez-Ramírez R, López-Bonilla OR, *et al.* Attention measurement of an autism spectrum disorder user using EEG Signals: A case study. *Mathe Comput Appl* 2022; 27(2): 21. <http://dx.doi.org/10.3390/mca27020021>
- [52] Pandian GSDB, Jain A, Raza Q, Sahu KK. Digital health interventions (DHI) for the treatment of attention deficit hyperactivity disorder (ADHD) in children - A comparative review of literature among various treatment and DHI. *Psychiatry Res* 2021; 297: 113742. <http://dx.doi.org/10.1016/j.psychres.2021.113742> PMID: 33515870
- [53] Low AM, Vangkilde S, le Sommer J, *et al.* Visual attention in adults with attention-deficit/hyperactivity disorder before and after stimulant treatment. *Psychol Med* 2019; 49(15): 2617-25. <http://dx.doi.org/10.1017/S0033291718003628> PMID: 30560740
- [54] Zabcikova M. Measurement of visual and auditory stimuli using EEG headset emotiv epoc. *MATEC Web Conf* 2019; 292: 01023. <http://dx.doi.org/10.1051/mateconf/201929201023>.
- [55] Ordikhani-Seyedlar M, Lebedev MA, Sorensen HBD, Puthusserypady S. Neurofeedback therapy for enhancing visual attention: State-of-the-art and challenges. *Front Neurosci* 2016; 10: 352. <http://dx.doi.org/10.3389/fnins.2016.00352> PMID: 27536212
- [56] Benedetti F, Catenacci Volpi N, Parisi L, Sartori G. Attention training with an easy-to-use brain computer interface. *Virtual, Augmented and Mixed Reality Applications of Virtual and Augmented Reality*. 236-47. http://dx.doi.org/10.1007/978-3-319-07464-1_22
- [57] Thomas KP, Vinod AP, Guan C. Design of an online EEG based neurofeedback game for enhancing attention and memory. 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC). Osaka, Japan. 2013; pp. 433-6. <http://dx.doi.org/10.1109/EMBC.2013.6609529>
- [58] Khan AA, Madendran RK, Thirunavukkarasu U, Faheem M. D²PAM : Epileptic seizures prediction using adversarial deep dual patch attention mechanism. *CAAI Trans Intell Technol* 2023; 8(3): 755-69. <http://dx.doi.org/10.1049/cit2.12261>
- [59] Belchior P, Marsiske M, Sisco SM, *et al.* Video game training to improve selective visual attention in older adults. *Comput Human Behav* 2013; 29(4): 1318-24. <http://dx.doi.org/10.1016/j.chb.2013.01.034> PMID: 24003265
- [60] Eivazi S, Bednarik R. Predicting problem-solving behavior and performance levels from visual attention data. *Proceedings of workshop on eye gaze in intelligent human machine interaction at IUI*. 9-16.
- [61] Salamati A, Hosseini SA, Haghgou H. Effectiveness of vestibular stimulation on visual attention in children with attention deficit hyperactivity disorder. *Archiv Rehabil* 2014; 15(3): 18-25.
- [62] Reimelt C, Wolff N, Hölling H, Mogwitz S, Ehrlich S, Roessner V. The underestimated role of refractive error (hyperopia, myopia, and astigmatism) and strabismus in children with ADHD. *J Atten Disord* 2021; 25(2): 235-44. <http://dx.doi.org/10.1177/1087054718808599> PMID: 30371126
- [63] Solé Puig M, Pérez Zapata L, Puigcerver L, *et al.* Attention-related eye vergence measured in children with attention deficit hyperactivity disorder. *PLoS One* 2015; 10(12): e0145281. <http://dx.doi.org/10.1371/journal.pone.0145281> PMID: 26694162
- [64] Karaca I, Demirkılınc Biler E, Palamar Onay M, Özbaran B, Üretmen Ö. Stereoacuity, fusional vergence amplitudes, and refractive errors prior to treatment in patients with attention-deficit hyperactivity disorder. *Turk J Ophthalmol* 2020; 50(1): 15-9. <http://dx.doi.org/10.4274/tjo.galenos.2019.17802> PMID: 32166943
- [65] Borsting E, Rouse M, Chu R. Measuring ADHD behaviors in children with symptomatic accommodative dysfunction or convergence insufficiency: A preliminary study. *Optometry* 2005; 76(10): 588-92. <http://dx.doi.org/10.1016/j.optm.2005.07.007> PMID: 16230274
- [66] Scheiman M, Mitchell GL, Cotter SA, *et al.* Convergence insufficiency treatment trial-attention and reading trial (CITT-ART): Design and methods. *Vis Dev Rehabil* 2015; 1(3): 214-28. PMID: 26942226
- [67] Redondo B, Vera J, Molina R, *et al.* Attention-deficit/hyperactivity disorder children exhibit an impaired accommodative response. *Graefes Arch Clin Exp Ophthalmol* 2018; 256(5): 1023-30. <http://dx.doi.org/10.1007/s00417-018-3948-2> PMID: 29569083

- [68] Findlay JM. Saccadic eye movement programming: Sensory and attentional factors. *Psychol Res* 2009; 73(2): 127-35.
- [69] CITT-ART Investigator Group. Treatment of symptomatic convergence insufficiency in children enrolled in the convergence insufficiency treatment trial-attention & reading trial: A randomized clinical trial. *Optom Vis Sci* 2019; 96(11): 825-35. <http://dx.doi.org/10.1097/OPX.0000000000001443>.
- [70] Gallaway M, Scheiman M, Mitchell GL. Vision therapy for post-concussion vision disorders. *Optom Vis Sci* 2017; 94(1): 68-73. <http://dx.doi.org/10.1097/OPX.0000000000000935> PMID: 27505624
- [71] Rucker JC, Phillips PH. Efferent vision therapy. *J Neuroophthalmol* 2018; 38(2): 230-6. <http://dx.doi.org/10.1097/WNO.0000000000000480> PMID: 28059865
- [72] Zohourian B, Shandiz JH, Riazi A, Khorasani AA, Yazdani N, Mostaeidi MT. Impact of vision therapy on eye-hand coordination skills in students with visual impairment. *J Ophthalmic Vis Res* 2018; 13(3): 301-6. http://dx.doi.org/10.4103/jovr.jovr_103_17 PMID: 30090187
- [73] Barrett BT. A critical evaluation of the evidence supporting the practice of behavioural vision therapy. *Ophthalmic Physiol Opt* 2009; 29(1): 4-25. <http://dx.doi.org/10.1111/j.1475-1313.2008.00607.x> PMID: 19154276
- [74] Jin P, Li X, Ma B, Guo H, Zhang Z, Mao L. Dynamic visual attention characteristics and their relationship to match performance in skilled basketball players. *PeerJ* 2020; 8: e9803. <http://dx.doi.org/10.7717/peerj.9803> PMID: 32879809
- [75] Alqarafi A, Ahmad Khan A, Kumar Mahendran R, Al-Sarem M, Albalwy F. Multi-scale GC-T2: Automated region of interest assisted skin cancer detection using multi-scale graph convolution and tri-movement based attention mechanism. *Biomed Signal Process Control* 2024; 95: 106313. <http://dx.doi.org/10.1016/j.bspc.2024.106313>
- [76] Khan AA, Mahendran RK, Perumal K, Faheem M. Dual-3DM 3-AD: Mixed transformer based semantic segmentation and triplet pre-processing for early multi-class Alzheimer's diagnosis. *IEEE Trans Neural Syst Rehabil Eng* 2024; 32: 696-707. <http://dx.doi.org/10.1109/TNSRE.2024.3357723> PMID: 38261494
- [77] Khan AA, Shahid MM, Bashir RN, *et al.* Detection of omicron caused pneumonia from radiology images using convolution neural network (CNN). *Comput Mater Continua* 2023; 74(2)
- [78] Kujur A, Raza Z, Khan AA, Wechtaisong C. Data complexity based evaluation of the model dependence of brain MRI images for classification of brain tumor and Alzheimer's disease. *IEEE Access* 2022; 10: 112117-33. <http://dx.doi.org/10.1109/ACCESS.2022.3216393>
- [79] Srinithya G. Effectiveness of vision therapy as an adjunct to occupational therapy in improving visual motor skills in learning disabled children. Doctoral thesis, JKK Muniraja Medical Research Foundation 2017.

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.