

# Time-Dependent Cortical Activation in Voluntary Muscle Contraction

Qi Yang<sup>1,4</sup>, Xiaofeng Wang<sup>3</sup>, Yin Fang<sup>1</sup>, Vlodek Siemionow<sup>1,2</sup>, Wanxiang Yao<sup>5</sup> and Guang H. Yue<sup>\*1,2,4</sup>

Departments of <sup>1</sup>Biomedical Engineering, <sup>2</sup>Physical Medicine and Rehabilitation and <sup>3</sup>Quantitative Health Sciences, The Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, USA

<sup>4</sup>Department of Chemical and Biomedical Engineering, Cleveland State University, Cleveland, Ohio 44115, USA

<sup>5</sup>Department of Health and Kinesiology, University of Texas at San Antonio, San Antonio, TX 78249, USA

**Abstract:** This study was to characterize dynamic source strength changes estimated from high-density scalp electroencephalogram (EEG) at different phases of a submaximal voluntary muscle contraction. Eight healthy volunteers performed isometric handgrip contractions of the right arm at 20% maximal intensity. Signals of the handgrip force, electromyography (EMG) from the finger flexor and extensor muscles and 64-channel EEG were acquired simultaneously. Sources of the EEG were analyzed at 19 time points across preparation, execution and sustaining phases of the handgrip. A 3-layer boundary element model (BEM) based on the MNI (Montréal Neurological Institute) brain MRI was used to overlay the sources. A distributed current density model, LORETA L1 norm method was applied to the data that had been processed by independent component analysis (ICA). Statistical analysis based on a mixed-effects polynomial regression model showed a significant and consistent time-dependent non-linear source strength change pattern in different phases of the handgrip. The source strength increased at the preparation phase, peaked at the force onset time and decreased in the sustaining phase. There was no significant difference in the changing pattern of the source strength among Brodmann's areas 1, 2, 3, 4, and 6. These results show, for the first time, a high time resolution increasing-and-decreasing pattern of activation among the sensorimotor regions with the highest activity occurs at the muscle activity onset. The similarity in the source strength time courses among the cortical centers examined suggests a synchronized parallel function in controlling the motor activity.

**Keywords:** EEG source, Current density reconstruction, Electroencephalography (EEG), Brain, Handgrip force, Voluntary muscle contraction.

## INTRODUCTION

The time course of brain activation during voluntary muscle contraction was first reported by the studies involving single-cell recordings in the brain of non-human primates [1-3]. These studies reported varying activation levels of motor cortical neurons among various phases (e.g., preparation vs. execution) during a motor control process. Although the observations provide valuable information describing single cell activities with time in animals in controlling a voluntary motor action, very little is known about activation time course of individual brain regions in humans in millisecond resolution.

Time courses of human cortical field activation are typically studied by neuroimaging involving relatively long-duration tasks. For example, alterations of activation of various brain areas were investigated during muscle fatigue lasting for many seconds using functional magnetic resonance imaging (fMRI) [4, 5] and positron emission tomography (PET) [6]. These studies demonstrate time- or fatigue-dependent alterations of cortical activities across a

relatively long time but cannot identify potential shifts of the activity level from one stage (e.g., planning) to another (e.g., execution) in a motor control process given the relatively poor temporal resolution of current-available neural imaging techniques. Although scalp electroencephalography (EEG) [7] or magnetoencephalography (MEG) [8] signals have millisecond resolution and are frequently recorded to describe control strategies of the brain, the source(s) of the signals or the involving cortical fields are usually unknown because of relatively poor spatial resolution of these methods. To improve spatial resolution and maintain high temporal resolution of scalp EEG or MEG data, several inverse solution methods to estimate the signal sources have been developed [9]. These inverse methods aim to de-blur the head volume conductor effect on the signal recorded from the scalp to identify the true electrophysiological sources in the cortex. With detailed temporal and spatial information, the inverse method could serve as a promising technique to explore activation time course of a cortical region or regions in movement control.

Commonly used inverse models include equivalent-current-dipole, distributed current density, beamforming, etc. Equivalent-current-dipole model relies on an ad hoc assumption of a reasonable number of dipoles, their known locations and/or orientations. Distributed current density model does not need prior knowledge of specific source number and

\*Address correspondence to this author at the Department of Biomedical Engineering, The Cleveland Clinic, 9500 Euclid Avenue, ND20, Cleveland, OH 44195, USA; Tel: (216) 4459336; Fax: (216) 444-9198; E-mail: yueg@ccf.org

locations. Among the several available current-density estimation techniques, the low-resolution electromagnetic tomography (LORETA) has been shown to have better source localization ability than other methods [10, 11]. The LORETA technique has been used in investigating abnormalities of P300 in schizophrenia [12, 13], analyzing widespread epileptiform activities in epilepsy [14, 15], and exploring cognitive function of the brain in healthy individuals [16].

The purpose of this study was to use LORETA source-localization technique to estimate sources of scalp EEG signals (64 channels) recorded before and during a handgrip motor task and determine the time-dependent source strength (activity level) changes across preparation, execution and sustaining phases of the task in different cortical sensorimotor areas. It was hypothesized that neural activity in the cortical regions would vary depending on the control phases of the handgrip task and the changes could be detected by the current density reconstruction technique.

## MATERIALS AND METHODOLOGY

### Subjects and Motor Task

Eight right-handed subjects (4 men and 4 woman, age =  $31 \pm 2$  yrs) participated in the study, none had neurological or other disorders at the time of study. The experimental procedures were approved by the Institutional Review Board at the Cleveland Clinic. All subjects gave informed consent prior to the participation.

Subjects were seated comfortably in an experimental chair in an electrically shielded data-recording room. During the experiment, subjects performed 20 isometric handgrip contractions of the right arm at 20% maximal voluntary contraction (MVC) level. An oscilloscope was placed in front of the subjects and the target was displayed on the screen. In each trial, they steadily increased the handgrip force to the target (20% MVC) level and matched the exerted force as closely as possible with the target for about 3 s.

Special precautions were taken to minimize noises in the data during the experiments. The subjects were required not to move the head and body, blink the eyes, bite the teeth, contract the facial/neck muscles, or tense the body while performing the low-intensity handgrips. Muscle fatigue was minimal by allowing sufficient rest between adjacent trials. The length of the inter-trial rest period was controlled by the subjects themselves, i.e., they started the next trial when they felt comfortable and completely rested. The inter-trial rest period was, on average about 20s.

### Data Recording

#### Force

The handgrip force was recorded by a pressure transducer (EPX-N1 250 PSI, Entran Devices, Inc., Fairfield, NJ) interfaced with distilled water in a bottle-like device through a nylon tube [17]. The output of the transducer was directed to the amplifier and then to an input channel of a Micro 1401 data acquisition system (version 3.05, Cambridge Electronic Design, Ltd., Cambridge, UK), which

transferred the voltage data to a laptop computer. The sampling rate for digitizing the force data was 100 Hz.

#### EEG

A 64-channel NeuroScan EEG system (version 4.2, NeuroScan, El Paso, Texas, USA) was used to acquire EEG signals from the scalp. A Quik-Cap elastic nylon cap that held 64 electrodes was placed on the scalp for EEG data recording. Conducting gel was injected into each electrode to connect the recording surface of the electrode with the scalp. Impedance was controlled below 10,000 ohms. One electrode was used to record the force, which was used for the purpose of synchronizing the EEG with muscle activities. All the EEG electrodes were referenced to the common linked earlobes and the signals were amplified ( $\times 75,000$ ), low-pass filtered (0.05-50 Hz), and digitized (2,000 sample/s).

#### Data Analysis

The force data were processed using house-coded programs within the Spike2 software package (Cambridge Electronic Design, Ltd., Cambridge, UK). The force onset time was set at the 10% handgrip MVC force rising from the baseline in each trial, which was used for the purpose of synchronizing the force, and EEG signals. The actual force was measured by averaging the data points in each trial when the force was steady (rising and falling portions were excluded) and then averaging across the trials before performing the group averaging.

The raw EEG data were visually inspected and trials with artifacts due to eye blinks or head movements were excluded (on average,  $2 \pm 2$  trials were removed in each subject). Both EEG data preprocessing and LORETA current density estimation were performed using Curry software package (version 4.5, Neuro Scan Labs, Virginia, USA).

#### Estimation of Current Densities Using LORETA

For each subject the EEG signals were aligned with the force onset and then averaged across trials. The averaged EEG data were baseline corrected using baseline data from -3000ms to -2500ms (time 0 indicates force onset). Subsequently, an independent component analysis (ICA) was applied to the data. Only the main components (signal-to-noise ratio  $> 1$ ) were chosen for the source reconstruction. A 3-layer (conductivities of the scalp and brain: 0.033S/m, and the skull: 0.0042S/m) triangle-node boundary element model (BEM) [18, 19] based on the MNI (Montréal Neurological Institute) brain MRI was used to overlay the sources. Distributed current density model (LORETA) with L1 norm method was applied to the ICA-preprocessed data. In addition, the sources were constrained to the reconstructed layer of the folded cortex with 6926 nodes using a rotating model [Curry user guide, 1999, 11].

#### Current Density Data Analysis - Post Processing

Source localization was analyzed at 21 different time points (-2200ms, -2100ms, -2000ms, -1000ms, -900ms, -800ms, -700ms, -600ms, -500ms, -400ms, -300ms, -200ms, -100ms, 0ms, 100ms, 200ms, 500ms, 800ms, 1000ms, 2000ms, 2500ms and 3000ms) throughout the planning,





**Table 1.** Analysis of Covariance Based on the Mixed-Effects Polynomial Regression Model. The "Time" Variable was Normalized into [0,1] in the Analysis. The p-Values Based on ANOVA F Tests are Presented for Global Fixed Effects Over all Subjects

		Estimate	Std Error	p-value
Intercept		1.10577	0.70643	0.0224
Time		-11.49543	6.81269	<.0001
Time^2		120.15607	21.14488	<.0001
Time^3		-191.55915	31.6776	<.0001
Time^4		84.28151	15.9606	<.0001
Area	Area1: Left M1	0	0	0.7255
	Area2: Right M1	-0.92206	0.85077	
	Area3: Left S1	0.05944	0.85077	
	Area4: Right S1	-0.50936	0.85077	
	Area5: PM&SMA	-0.70075	0.85077	

Normalized Current Density

Time (ms)

**Fig. (3).** The original data and fitted curves for each area within each subject from the mixed-effects polynomial regression model. Each column represents the case of one subject and each row represents the case of one area. Area 1, area2, area3, area 4, and area5 denoted the left M1, right M1, left S1, right S1, and PM and SMA, respectively. For example, the sub-plot subject5/area3 denotes the case of subject 5 with area3: Left S1.

**Time-Dependent Source Strength Variation in Controlling a Voluntary Handgrip**

Our results indicated that from as early as 1 s before the force onset the source strength had elevated in cortical regions examined (Fig. 1, 2 and 3) and peaked near the time of force onset. Cortical activation at early times may be more related to attention and general preparation of a voluntary motor action [23]. At early stages of motor preparation,

perhaps only a limited number of neurons are involved in the

task, and the activation level might not critically depend on the force exerted [24-29]. As the time approaches motor task initiation, more activities related to specific programming and execution of the task occur and this may require progressively more cortical neurons to participate and increase their activation levels. This postulation is supported by the almost linear increases of the source strength from -1000ms

to the force onset (time 0, Fig. 2). The observation of the source strength at the highest level around the force onset could be contributed by participation of neurons that executed the handgrip and neurons whose role involved in modulating the grip force. Cheney [30] suggested that about 100ms before the EMG onset, force-magnitude modulation, known as increasing discharge rate and/or recruitment of more neurons, might have been established. About 100ms or more after the EMG onset, this modulation is further adjusted so that the cortical activity increases according to the level of force generated by the muscle. In summary, the high levels of cortical activation during the time window of  $\pm 100$ ms movement onset are highly likely to be related to specific motor task programming, execution, force modulation, and perhaps as well as multi-level sensory feedback reaching to the motor cortices [30].

During the later phase of the holding period ( $\sim 200$ ms after the force onset), the source strength began to decline in all the analyzed areas (Fig. 2, Fig. 3). The reduction in the cortical activity during the holding phase could be a result of ceasing activation of the neurons involved in motor action programming and execution. Many cortical cells increase discharge rate before the force onset but most of these neurons were silent during the period of holding the force. For example, more than 60% of the recorded cells in monkey SMA and cingulate cortex were phasically active before onset of finger force but only about 10% of the neurons were tonically active during the holding phase of the force task [31]. In addition, motor neurons in the spinal cord adapt to lower firing rate after initial high-frequency activation [32] and this adaptation could also occur in cortical neurons [33], which may as well explain weakening of the source strength during later phase of the force holding.

### Similar Source Strength Variation Time Courses among the Cortical Regions

All the examined regions showed time-dependent source strength changes during the control process of the muscle contraction. The left M1 and S1 exhibited stronger current density (CD) during almost the entire time course (Fig. 2), which was expected because the handgrip occurred on the right side. An interesting observation was that although the level of the CD seemed to differ among the cortical areas, the pattern of the source strength variation in the time course was very similar, that is all the areas exhibited a reversed "V-shape" CD time course (Fig. 2). This was confirmed by the statistical F-test that all terms of polynomial coefficients for the nonlinear current density time course were significant while the ANOVA test for the factor of "area" was not. Previous animal studies involving single-cell recording reported that the M1 is primarily for controlling movement execution variables, such as the amount of force to be exerted by the muscle [1, 30] and direction of the force production [34]. The M1 also encodes higher functions, such as movement programming and preparation [35]. The S1 is known to participate in handgrip force control [36] and provide sensory information to M1 during voluntary motor actions [37]. The SMA has been shown to be important in programming sequential finger movements [38], storing information necessary for the orderly performance of multiple movements of the arm [39], and bimanual coordination

[40, 41]. The PM is considered to contribute to the selection of motor actions on the basis of visual cues [42, 43]. The

monotonic relationship between cell activity and muscle output holds for different areas in the motor cortex in a series of single-cell animal studies with slightly different experiment paradigms [1, 44-46]. Specifically, the M1 and PM cortices share the same force-firing rate pattern [46]. Two populations of neurons (phasic-tonic firing and tonic decreasing firing) in the M1 and PM cortex have been found to be related, either positively or negatively, to force output in a more recent study [47]. The relationship between the pyramidal tract neurons and force output was monotonic over only a small portion of the force range [45]. In summary, all the examined areas play important motor function, which is underscored by their contributions to the corticospinal tract (CST), a fundamental pathway for voluntary actions of extremity muscles (one third of the CST is contributed by the M1, one third by the PM and SMA, and the remaining one third by the S1 [48]).

One limitation of the single-cell recording in animal models, however, is the difficulty to investigate activities of multiple cells in multiple cortical regions concurrently. Thus, it is not well known, from the animal data, the integrated activation patterns of multiple brain areas. This limitation can be overcome by modern neuroimaging techniques and abundant observations have suggested high degrees similarities in activation patterns of many sensorimotor function-related cortical areas. For example, neuroimaging data have shown a proportional relationship between the levels of muscle and activation, and this relation is similar among many cortical regions [6, 36, 49, 50]. A number of studies have reported muscle fatigue-related brain activation alterations and a prominent finding by these studies was that the cortical signal changing pattern was very much alike among many sensorimotor regions [4, 5, 17]. The human neuroimaging data support the observation of similar source strength time courses among the cortical areas in the current study and suggest synchronized parallel functions among multiple sensorimotor centers.

Compared with previous findings of EEG source signals generated during higher levels of handgrip contractions, the relative source strength (CD) of the SMA and PM (relative to CD of M1 and S1) was greater for the 20% MVC of the current study than relative source strength of the SMA and PM at higher levels of handgrip [51]. It is not clear why this was so but it could be that fewer cells in the primary sensorimotor areas need to be activated to plan, execute and maintain a low level of muscle contraction but the level of activation in the SMA and PM may not change as much across degrees of muscle contraction.

In conclusion, the results show that the level of cortical activity in controlling a voluntary motor action is time-dependent with the highest activity occurs around the action onset. The low-to-high-to-low source strength pattern from the preparation to execution and to holding phases is similar among several major sensorimotor areas, suggesting synchronized parallel functions among them. The findings also indicate that the current density reconstruction technique based on surface EEG recorded during voluntary motor

activities has adequate time and spatial resolution to detect the time-dependent source strength in various brain regions.

## ACKNOWLEDGEMENTS

This work was supported by research funds of Department of Physical Medicine and Rehabilitation at the Cleveland Clinic

## REFERENCES

- [1] Evarts EV. Relation of pyramidal tract activity to force exerted during voluntary movement. *J Neurophysiol* 1968; 31: 14-27.
- [2] Conrad B, Wiesendanger M, Matsunami K, Brooks VB. Precentral unit activity related to control of arm movements. *Exp Brain Res* 1977; 29: 85-95.
- [3] Fetz EE, Cheney PD. Postspike facilitation of forelimb muscle activity by primate corticomotoneuronal cells. *J Neurophysiol* 1980; 44: 751-72.
- [4] Liu JZ, Shan ZY, Zhang LD, Sahgal V, Brown RW, Yue GH. Human brain activation during sustained and intermittent submaximal fatigue muscle contractions: an fMRI study. *J Neurophysiol* 2003; 90: 300-12.
- [5] van Duinen H, Renken R, Maurits N, Zijdwend I. Effects of motor fatigue on human brain activity, an fMRI study. *Neuroimage* 2007; 35: 1438-49.
- [6] Dettmers C, Fink GR, Lemon RN, et al. Relation between cerebral activity and force in the motor areas of the human brain. *J Neurophysiol* 1995; 74: 802-15.
- [7] Siemionow V, Yue GH, Ranganathan VK, Liu JZ, Sahgal V. Relationship between motor activity-related cortical potential and voluntary muscle activation. *Exp Brain Res* 2000; 133: 303-11.
- [8] Tecchio F, Zappasodi F, Melgari JM, Porcaro C, Cassetta E, Rossini PM. Sensory-motor interaction in primary hand cortical areas: a magnetoencephalography assessment. *Neuroscience* 2006; 141(1): 533-42.
- [9] Michel CM, Murray MM, Lantz G, Gonzalez S, Spinelli L, Grave de Peralta R. EEG source imaging. *Clin Neurophysiol* 2004; 115: 2195-222.
- [10] Babiloni F, Babiloni C, Locche L, Cincotti F, Rossini PM, Carducci F. High-resolution electro-encephalogram: source estimates of Laplacian-transformed somatosensory-evoked potentials using a realistic subject head model constructed from magnetic resonance images. *Med Biol Eng Comput* 2000; 38: 512-19.
- [11] Yao J, Dewald JP. Evaluation of different cortical source localization methods using simulated and experimental EEG data. *Neuroimage* 2005; 25: 369-82.
- [12] Winterer G, Mulert C, Mientus S, et al. P300 and LORETA: comparison of normal subjects and schizophrenic patients. *Brain Topogr* 2001; 13: 299-13.
- [13] Pae JS, Kwon JS, Youn T, et al. LORETA imaging of P300 in schizophrenia with individual MRI and 128-channel EEG. *Neuroimage* 2003; 20: 1552-60.
- [14] Lantz G, Michel CM, Pascual-Marqui RD, et al. Extracranial localization of intracranial interictal epileptiform activity using LORETA (low resolution electromagnetic tomography). *Electroencephalogr Clin Neurophysiol* 1997; 102: 414-22.
- [15] Worrell GA, Lagerlund TD, Sharbrough FW, et al. Localization of the epileptic focus by low-resolution electromagnetic tomography in patients with a lesion demonstrated by MRI. *Brain Topogr* 2000; 12: 273-82.
- [16] Thut G, Hauert CA, Morand S, Seeck M, Landis T, Michel C. Evidence for interhemispheric motor-level transfer in a simple reaction time task: an EEG study. *Exp Brain Res* 1999; 128: 256-61.
- [17] Liu JZ, Dai TH, Sahgal V, Brown RW, Yue GH. Non-linear cortical modulation of muscle fatigue: A functional MRI study. *Brain Research* 2002; 957: 320-29.
- [18] Fuchs M, Drenckhahn R, Wischmann HA, Wagner M. An improved boundary element method for realistic volume-conductor modeling. *IEEE Trans Biomed Eng* 1998; 45: 980-97.
- [19] Fuchs M, Kastner J, Wagner M, Hawes S, Ebersole JS. A standardized boundary element method volume conductor model. *Clin Neurophysiol* 2002; 113: 702-12.
- [20] Hedeker D, Gibbons RD. Longitudinal data analysis. Chapter 5. John Wiley & Sons 2006.
- [21] Davis CS. Statistical methods for the analysis of repeated measurements. New York: Springer 2002.
- [22] Verbeke G, Molenberghs G. Linear mixed models for longitudinal data. New York: Springer 2000.
- [23] Ball T, Schreiber A, Feige B, Wagner M, Lucking CH, Kristeva-Feige R. The role of higher-order motor areas in voluntary movement as revealed by high-resolution EEG and fMRI. *Neuroimage* 1999; 10: 682-94.
- [24] Buys EJ, Lemon RN, Mantel GW, Muir RB. Selective facilitation of different hand muscles by single corticospinal neurones in the conscious monkey. *J Physiol* 1986; 381: 529-49.
- [25] Ashe J. Force and the motor cortex. *Behav Brain Res* 1997; 87: 255-69.
- [26] Shen L, Alexander GE. Neural correlates of a spatial sensory-to-motor transformation in primary motor cortex. *J Neurophysiol* 1997; 77: 1171-94.
- [27] Sakata H, Taira M, Kusunoki M, Murata A, Tanaka Y. The TINS Lecture. The parietal association cortex in depth perception and visual control of hand action. *Trends Neurosci* 1997; 20: 350-7.
- [28] Murata A, Gallese V, Luppino G, Kaseda M, Sakata H. Selectivity for the shape, size, and orientation of objects for grasping in neurons of monkey parietal area AIP. *J Neurophysiol* 2000; 83: 2580-601.
- [29] Morrow MM, Miller LE. Prediction of muscle activity by populations of sequentially recorded primary motor cortex neurons. *J Neurophysiol* 2003; 89: 2279-88.
- [30] Cheney PD. Role of cerebral cortex in voluntary movements. *A review Phys Ther* 1985; 65: 624-35.
- [31] Cadoret G, Smith AM. Comparison of the neuronal activity in the SMA and the ventral cingulate cortex during prehension in the monkey. *J Neurophysiol* 1997; 77: 153-66.
- [32] Kernell D, Monster AW. Motoneurone properties and motor fatigue: an intracellular study of gastrocnemius motoneurons of the cat. *Exp Brain Res* 1982; 46: 197-204.
- [33] Hepp-Reymond M-C, Wannier TMJ, Maier MA, Rufener EA. Sensorimotor cortical control of isometric force in the monkey. *Progress in Brain Res* 1989; 80: 451-63.
- [34] Georgopoulos AP, Schwartz AB, Kettner RE. Neuronal population coding of movement direction. *Science* 1986; 233: 1416-9.
- [35] Michelon P, Vettel JM, Zacks JM. Lateral somatotopic organization during imagined and prepared movements. *J Neurophysiol* 2006; 95: 811-22.
- [36] Dai TH, Liu JZ, Sahgal V, Brown RW, Yue GH. Relationship between muscle output and functional MRI-measured brain activation. *Exp Brain Res* 2001; 140: 290-300.
- [37] Berne RM, Levy MN, Koeppen BM, Stanton BA. *Physiology-5<sup>th</sup> edition*. Mosby
- [38] Roland PE, Larsen B, Lassen NA, Skinhoj E. Supplementary motor area and other cortical areas in organization of voluntary movements in man. *J Neurophysiol* 1980; 43: 118-36.
- [39] Tanji J, Shima K. Role for supplementary motor area cells in planning several movements ahead. *Nature* 1994; 371: 413-6.
- [40] Brinkman C. Lesions in supplementary motor area interfere with a monkey's performance of a bimanual coordination task. *Neurosci Lett* 2003; 27: 267-70.
- [41] Brinkman C. Supplementary motor area of the monkey's cerebral cortex: short- and long-term deficits after unilateral ablation and the effects of subsequent callosal section. *J Neurosci* 1984; 4: 918-29.
- [42] Halsband U, Passingham RE. Premotor cortex and the conditions for movement in monkeys (*Macaca fascicularis*). *Behav Brain Res* 1985; 18: 269-77.
- [43] Passingham RE. Cues for movement in monkeys (*Macaca mulatta*) with lesions in premotor cortex. *Behav Neurosci* 1986; 100: 695-703.
- [44] Hepp-Reymond MC, Wyss UR and Anner R. Neuronal coding of static force in the primate motor cortex. *J Physiol (Paris)* 1978; 74: 287-91.
- [45] Cheney PD, Fetz EE. Functional classes of primate corticomotoneuronal cells and their relation to active force. *J Neurophysiol* 1980; 44: 773-91.

- [46] Hepp-Reymond M, Kirkpatrick-Tanner M, Gabernet L, Qi HX, Weber B. Context-dependent force coding in motor and premotor cortical areas. *Exp Brain Res* 1999; 128: 123-33.
- [47] Hepp-Reymond MC, Hüsler EJ, Maier MA, Qi HX. Force-related neuronal activity in two regions of the primate ventral premotor cortex. *Can J Physiol Pharmacol* 1994; 72: 571-9.
- [48] Ghez C, Krakauer J. The Organization of Movement. In: E.R. Kandel, J.H. Schwartz and T.M. Jessell (Eds.), *Principles of Neural Science* (4th ed). McGraw-Hill, New York, 2000; 653-73.
- [49] Thickbroom GW, Phillips BA, Morris I, Byrnes ML, Mastaglia FL. Isometric force-related activity in sensorimotor cortex measured with functional MRI. *Exp Brain Res* 1998; 121: 59-64.
- [50] Cramer SC, Weisskoff RM, Schaechter JD, *et al.* Motor cortex activation is related to force of squeezing. *Hum Brain Mapp* 2002; 16: 197-205.
- [51] Wang XF, Yang Q, Fan Z, Sun CK, Yue GH. Assessing time-dependent association between scalp EEG and muscle activation: A functional random-effects model approach. *J Neurosci Methods* 2009; 177: 232-40.

---

Received: November 06, 2010

Revised: May 10, 2011

Accepted: June 27, 2011

© Yang *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.